

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/52649 A1

(51) International Patent Classification⁷: A01N 1/02,
C12N 5/00, 5/02, 7/00, 15/63, 15/86, C12P 21/04, 21/06

69 Peter Courtts Circle, Stanford, CA 94305 (US). **WEISS-
MAN, Irving, L.**; 4147 Jefferson Avenue, Redwood City,
CA 94061 (US).

(21) International Application Number: PCT/US01/01459

(22) International Filing Date: 17 January 2001 (17.01.2001)

(74) Agent: **SHERWOOD, Pamela, J.**; Bozicevic, Field &
Francis LLP, Suite 200, 200 Middlefield Road, Menlo Park,
CA 94025 (US).

(25) Filing Language: English

(81) Designated States (*national*): AU, CA.

(26) Publication Language: English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(30) Priority Data:
60/176,786 18 January 2000 (18.01.2000) US

(71) Applicant: **THE BOARD OF TRUSTEES OF THE LE-
LAND STANFORD JUNIOR UNIVERSITY** [US/US];
Suite 350, 900 Welch Road, Palo Alto, CA 94304 (US).

Published:
— with international search report

(72) Inventors: **REYA, Tannishtha**; 777 W. Middlefield Road
#200, Mountain View, CA 94043 (US). **NUSSE, Roeland**;

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.



BEST AVAILABLE COPY

WO 01/52649 A1

(54) Title: EXPANSION OF STEM AND PROGENITOR CELLS BY BETA-CATENIN

(57) Abstract: Mammalian progenitor or stem cells are expanded *in vitro* by increasing the levels of β -catenin in the cell. The ex-
panded cells substantially maintain their original phenotype including the ability to give rise to multiple types of differentiated cells.
The intracellular levels of β -catenin may be manipulated by providing exogenous β -catenin protein to the cell, or by introduction
into the cell of a genetic construct encoding β -catenin. The β -catenin may be a wild-type or stabilized mutant form of the protein.
Preferably the long term cell culture medium substantially lacks stromal cells and cytokines.

EXPANSION OF STEM AND PROGENITOR CELLS BY BETA-CATENIN

INTRODUCTION

Beta-catenin is a pivotal player in the signaling pathway initiated by Wnt proteins, which are mediators of several developmental processes. Beta-catenin activity is controlled by a large number of binding partners that affect the stability and the localization of beta-catenin, and it is thereby able to participate in such varying processes as gene expression and cell adhesion. Activating mutations in beta-catenin and in components regulating its stability have been found to contribute to upregulation of cell proliferation.

The β -catenin protein becomes stabilized in response to Wnt/Wg, moves to the nucleus and forms complexes with the LEF1/TCF transcription factors to regulate gene expression. The level of cytosolic β -catenin is determined by its interaction with a number of proteins including those in a multiprotein complex of Axin, GSK-3 β , APC and other proteins. The mechanism by which the Wnt signal is transmitted to this complex is unclear but it involves interaction of Wnt with its receptors, which are members of Frizzled family of seven transmembrane proteins. The stabilization of β -catenin stimulates the expression of genes including c-myc, c-jun, fra-1, and cyclin D1. This pathway is negatively regulated by Axin.

Beta-catenin is also an adherens junction protein. Adherens junctions are critical for the establishment and maintenance of epithelial layers, such as those lining organ surfaces.

AJs mediate adhesion between cells, communicate a signal that neighboring cells are present, and anchor the actin cytoskeleton. In serving these roles, AJs regulate normal cell growth and behavior. At several stages of embryogenesis, wound healing, and tumor cell metastasis, cells form and leave epithelia. This process, which involves the disruption and reestablishment of epithelial cell-cell contacts, may be regulated by the disassembly and assembly of AJs. AJs may also function in the transmission of the 'contact inhibition' signal, which instructs cells to stop dividing once an epithelial sheet is complete.

For many purposes, there is an interest in being able to expand stem and progenitor cells in culture. However, it is not simply a matter of maintaining cell viability for the stem cells, but also of ensuring that the stem cells increase in numbers without losing their distinctive phenotype. Current protocols for the *in vitro* culture of hematopoietic stem cells generally require one or a cocktail of cytokines, such as c-kit ligand (stem cell growth factor), flt-3, thrombopoietin, IL-6, *etc.* While a substantial increase in cell number can be obtained with such cultures, they do not provide for expanded number of cells that retain a capacity for long term repopulation of all hematopoietic lineages. See Domen and Weissman (1999) *Mol Med Today* 5(5):201-8; or Ziegler and Kanz (1998) *Curr Opin Hematol* 5(6):434-40.

Stem cells have also been grown in co-culture with stromal cells. However, it is particularly desirable to expand stem cells in a culture of known composition, rather than relying upon the presence of other cells for their maintenance.

There continues to be a strong demand for improvements in the *in vitro* culture of stem cells and progenitor cells. The present invention addresses this need.

SUMMARY OF THE INVENTION

Methods are provided for the expansion of progenitor or stem cells *in vitro*, whereby the cells retain their pluripotential phenotype after expansion. The intracellular level of β -catenin is increased in the cells in culture, either by providing exogenous β -catenin protein to the cell, or by introduction into the cell of a genetic construct encoding β -catenin. The β -catenin may be a wild type protein appropriate for the species from which the cells are derived, or preferably, a stabilized mutant form of the protein. The alteration in cellular levels of β -catenin provide for increased number of cells in cycle, and leads to cultures that containing proliferating cells that maintain an undifferentiated phenotype *in vitro*. The expanded cell populations are useful as a source of stem cells, e.g. to reconstitute function in a host that is deficient in a particular cell lineage or lineages. In one embodiment of the invention, the target cells are hematopoietic stem cells, which may be used in transplantation to restore hematopoietic function to autologous or allogeneic recipients.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Activated beta-catenin retrovirus induces increased growth of stem cells. Stem cells infected with control or beta-catenin-GFP retrovirus were sorted and cultured on 96 well plates for two days in the presence or absence of steel factor, and cell numbers were counted at the end of the culture period.

Figure 2. Stem cells infected with beta-catenin retain many stem cell markers in long term culture. Beta-catenin infected stem cell spheres were harvested from long term cultures at 5 weeks, trypsinized and allowed to express their surface proteins for 12 hours. Subsequently they were harvested and stained with antibodies to Thy1.1, Sca1, c-kit, and lineage antigens (B220, Mac-1, Gr-1, Ter119, CD5, CD3, CD8/4).

Figure 3. Stem cells infected with beta-catenin have the ability to give rise to multiple lineages when transplanted. 100,000 beta-catenin infected stem cells were harvested from long term cultures at 7 weeks, trypsinized and injected into lethally irradiated (950 Rads) allotype marked recipients along with 300,000 rescuing bone marrow cells from the host. Analysis of reconstitution along various lineages was carried out at 4 weeks after transplantation.

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

Mammalian progenitor or stem cells are expanded *in vitro* by increasing the levels of β -catenin in the cell. The intracellular levels of β -catenin may be manipulated by providing exogenous β -catenin protein to the cell, or by introduction into the cell of a genetic construct encoding β -catenin. The β -catenin may be a wild-type or stabilized mutant form of the protein. Preferably the long term cell culture medium substantially lacks stromal cells and cytokines. Cultures that provide stem cell activity can be obtained for at least three weeks, frequently six weeks and can be eight weeks or more. The culture media that are employed are conventional media for the growth of mammalian cells, optionally in the absence of serum using only defined protein factors. In the absence of the β -catenin, the medium is inefficient at maintaining growth of the undifferentiated cells.

In the first few days of culture, the expansion of stem/progenitor cells is limited, usually the number of phenotypic stem/progenitor cells is maintained, or slightly increased. After 2 to 3 weeks in the subject culture conditions, there is a substantial proliferation of cells having the desired phenotype, where the number of cells having a functional stem/progenitor cell phenotype is expanded.

DEFINITIONS

It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, animal species or genera, and reagents described, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

As used herein the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the culture" includes reference to one or more cultures and equivalents thereof known to those skilled in the art, and so forth. All technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention belongs unless clearly indicated otherwise.

β -catenin: The term β -catenin, as used herein, is intended to refer to both wild-type and stabilized forms of the β -catenin protein, and to fusion proteins and derivatives thereof. Usually the protein will be of mammalian origin, although the protein from other species may find use. The protein is conserved between species, for example the human sequence is active in mouse cells. The sequences of many β -catenin proteins are publicly known. For convenience, the sequences of the human and mouse homologs of this protein are provided

in the sequence listing, as SEQ ID NO:1; and SEQ ID NO:2, respectively. In one embodiment of the invention, a stabilized form of beta-catenin is used.

The ubiquitin-dependent proteolysis system is involved in the regulation of beta-catenin turnover. Beta-catenin becomes stabilized when proteasome-mediated proteolysis is inhibited and this leads to the accumulation of multi-ubiquitinated forms of beta-catenin (Aberle et al. (1997) EMBO J 16(13):3797-804). Substitution of the serine residues in the glycogen synthase kinase 3 β (GSK3beta) phosphorylation consensus motif of beta-catenin inhibits ubiquitination and results in stabilization of the protein. Examples of stabilized β -catenins include those with the amino acid changes D32Y; D32G; S33F; S33Y; G34E; S37C; S37F; T41I; S45Y; and deletion of AA 1-173. A number of publications describe stabilized β -catenin mutations. For example, see Morin et al. (1997) Science 275(5307):1787-90; Palacios et al. (1998) Cancer Res 58(7):1344-7; Muller et al. (1998) Genes Chromosomes Cancer 22(1):37-41; Miyoshi et al. (1998) Cancer Res 58(12):2524-7; Zurawel et al. (1998) Cancer Res. 58, 896-899; Voeller et al. (1998) Cancer Res. 58, 2520-2526; *etc.*

The sequence of the beta-catenin polypeptide may be altered in various ways known in the art to generate targeted changes in sequence. The polypeptide will usually be substantially similar to the sequences provided herein, i.e. will differ by at least one amino acid, and may differ by at least two but not more than about ten amino acids. Deletions may further include larger changes, such as deletions of a domain or exon, providing for active peptide fragments of the protein. Other modifications of interest include tagging, e.g. with the FLAG system, HA, green fluorescent protein, etc. Such alterations may be used to alter properties of the protein, by affecting the stability, specificity, etc. The protein may be joined to a wide variety of other oligopeptides or proteins for a variety of purposes, particular for facilitating transport across membranes.

Techniques for *in vitro* mutagenesis of cloned genes are known. Examples of protocols for scanning mutations may be found in Gustin et al., Biotechniques 14:22 (1993); Barany, Gene 37:111-23 (1985); Colicelli et al., Mol Gen Genet 199:537-9 (1985); and Prentki et al., Gene 29:303-13 (1984). Methods for site specific mutagenesis can be found in Sambrook et al., Molecular Cloning: A Laboratory Manual, CSH Press 1989, pp. 15.3-15.108; Weiner et al., Gene 126:35-41 (1993); Sayers et al., Biotechniques 13:592-6 (1992); Jones and Winistorfer, Biotechniques 12:528-30 (1992); Barton et al., Nucleic Acids Res 18:7349-55 (1990); Marotti and Tomich, Gene Anal Tech 6:67-70 (1989); and Zhu Anal Biochem 177:120-4 (1989).

Expression construct: In one embodiment of the invention, the beta-catenin is delivered to the targeted stem or progenitor cells by introduction of an exogenous nucleic acid expression vector into the cells. Many vectors useful for transferring exogenous genes

into target mammalian cells are available. The vectors may be episomal, e.g. plasmids, virus derived vectors such cytomegalovirus, adenovirus, etc., or may be integrated into the target cell genome, through homologous recombination or random integration, e.g. retrovirus derived vectors such MMLV, HIV-1, ALV, etc.

5 Retrovirus based vectors have been shown to be particularly useful when the target cells are hematopoietic stem cells. For example, see Baum *et al.* (1996) J Hematother 5(4):323-9; Schwarzenberger *et al.* (1996) Blood 87:472-478; Nolte *et al.* (1996) P.N.A.S. 93:2414-2419; and Maze *et al.* (1996) P.N.A.S. 93:206-210. Lentivirus vectors have also been described for use with hematopoietic stem cells, for example see Mochizuki *et al.* 10 (1998) J Virol 72(11):8873-83. The use of adenovirus based vectors with hematopoietic cells has also been published, see Ogniben and Haas (1998) Recent Results Cancer Res 144:86-92.

Various techniques known in the art may be used to transfect the target cells, e.g. electroporation, calcium precipitated DNA, fusion, transfection, lipofection and the like. The 15 particular manner in which the DNA is introduced is not critical to the practice of the invention.

Combinations of retroviruses and an appropriate packaging line may be used, where the capsid proteins will be functional for infecting the target cells. Usually, the cells and virus will be incubated for at least about 24 hours in the culture medium. Commonly used 20 retroviral vectors are "defective", i.e. unable to produce viral proteins required for productive infection. Replication of the vector requires growth in the packaging cell line.

The host cell specificity of the retrovirus is determined by the envelope protein, env (p120). The envelope protein is provided by the packaging cell line. Envelope proteins are of at least three types, ecotropic, amphotropic and xenotropic. Retroviruses packaged with 25 ecotropic envelope protein, e.g. MMLV, are capable of infecting most murine and rat cell types. Ecotropic packaging cell lines include BOSC23 (Pear *et al.* (1993) P.N.A.S. 90:8392-8396). Retroviruses bearing amphotropic envelope protein, e.g. 4070A (Danos *et al.*, *supra.*), are capable of infecting most mammalian cell types, including human, dog and mouse. Amphotropic packaging cell lines include PA12 (Miller *et al.* (1985) Mol. Cell. Biol. 5:431- 30 437); PA317 (Miller *et al.* (1986) Mol. Cell. Biol. 6:2895-2902) GRIP (Danos *et al.* (1988) PNAS 85:6460-6464). Retroviruses packaged with xenotropic envelope protein, e.g. AKR env, are capable of infecting most mammalian cell types, except murine cells.

The sequences at the 5' and 3' termini of the retrovirus are long terminal repeats (LTR). A number of LTR sequences are known in the art and may be used, including the 35 MMLV-LTR; HIV-LTR; AKR-LTR; FIV-LTR; ALV-LTR; etc. Specific sequences may be accessed through public databases. Various modifications of the native LTR sequences are also known. The 5' LTR acts as a strong promoter, driving transcription of the β -catenin

gene after integration into a target cell genome. For some uses, however, it is desirable to have a regulatable promoter driving expression. Where such a promoter is included, the promoter function of the LTR will be inactivated. This is accomplished by a deletion of the U3 region in the 3' LTR, including the enhancer repeats and promoter, that is sufficient to inactivate the promoter function. After integration into a target cell genome, there is a rearrangement of the 5' and 3' LTR, resulting in a transcriptionally defective provirus, termed a "self-inactivating vector".

Suitable inducible promoters are activated in a desired target cell type, either the transfected cell, or progeny thereof. By transcriptional activation, it is intended that transcription will be increased above basal levels in the target cell by at least about 100 fold, more usually by at least about 1000 fold. Various promoters are known that are induced in hematopoietic cell types, e.g. IL-2 promoter in T cells, immunoglobulin promoter in B cells, etc.

Preferred genetic constructs are those that can be removed from the target cells after expansion. This can be accomplished by the use of a transient vector system, or by including a heterologous recombination site that flanks the beta-catenin coding sequence. In this manner, after expansion the construct can be removed prior to use of the expanded cell population. Preferably a detectable marker, e.g. green fluorescent protein, luciferase, cell surface proteins suitable for antibody selection methods, etc. is included in the expression vector, such that after deletion of the construct the cells can be readily isolated that lack the exogenous beta-catenin.

The term "heterologous recombination site" is meant to encompass any introduced genetic sequence that facilitates site-specific recombination. In general, such sites facilitate recombination by interaction of a specific enzyme with two such sites. Exemplary heterologous recombination sites include, but are not necessarily limited to, *lox* sequences with recombination mediated by Cre enzyme; *frt* sequences (Golic et al. (1989) *Cell* 59:499-509; O'Gorman et al. (1991) *Science* 251:1351-5; recombination mediated by the FLP recombinase), the recognition sequences for the pSR1 recombinase of *Zygosaccharomyces rouxii* (Matsuzaki et al. (1990) *J. Bacteriol.* 172:610-8), and the like.

Sequences encoding *lox* sites are of particular interest for use in the present invention. A *lox* site is a nucleotide sequence at which the gene product of the *cre* gene, referred to herein as "Cre," catalyzes site-specific recombination. A particularly preferred *lox* site is a *loxP* site. The sequence of *loxP*, which is 34 bp in length, is known and can be produced synthetically or can be isolated from bacteriophage P1 by methods known in the art (see, e.g. Hoess et al. (1982) *Proc. Natl. Acad. Sci. USA* 79:3398). The *loxP* site is composed of two 13 bp inverted repeats separated by an 8 bp spacer region. Other suitable

lox sites include *loxB*, *loxL*, and *loxR*, which can be isolated from *E. coli* (Hoess et al. (1982) *Proc. Natl. Acad. Sci. USA* 22:3398).

In an alternative method, expression vectors that provide for the transient expression in mammalian cells may be used. In general, transient expression involves the use of an expression vector that is able to replicate efficiently in a host cell, such that the host cell accumulates many copies of the expression vector and, in turn, synthesizes high levels of a desired polypeptide encoded by the expression vector. Transient expression systems, comprising a suitable expression vector and a host cell, allow for the convenient short term expansion of cells, but do not affect the long term genotype of the cell.

Translocation modified β -catenin: In some cases it is desirable to provide exogenous β -catenin protein, rather than transducing the cells with an expression construct. The beta-catenin may be added to the culture medium at high levels. Preferably the beta-catenin is modified so as to increase its transport into the cells.

In one embodiment of the invention, tat protein is used to deliver beta-catenin. The preferred transport polypeptides are characterized by the presence of the tat basic region amino acid sequence (amino acids 49-57 of naturally-occurring tat protein); the absence of the tat cysteine-rich region amino acid sequence (amino acids 22-36 of naturally-occurring tat protein) and the absence of the tat exon 2-encoded carboxy-terminal domain (amino acids 73-86 of naturally-occurring tat protein). Transport polypeptides are attached to beta-catenin by chemical cross-linking or by genetic fusion, where the beta-catenin moiety may be a wild-type or stabilized form. A unique terminal cysteine residue is a preferred means of chemical cross-linking.

Stem cell: The term stem cell is used herein to refer to a mammalian cell that has the ability both to self-renew, and to generate differentiated progeny (see Morrison *et al.* (1997) *Cell* 88:287-298). Generally, stem cells also have one or more of the following properties: an ability to undergo asynchronous, or symmetric replication, that is where the two daughter cells after division can have different phenotypes; extensive self-renewal capacity; capacity for existence in a mitotically quiescent form; and clonal regeneration of all the tissue in which they exist, for example the ability of hematopoietic stem cells to reconstitute all hematopoietic lineages. "Progenitor cells" differ from stem cells in that they typically do not have the extensive self-renewal capacity, and often can only regenerate a subset of the lineages in the tissue from which they derive, for example only lymphoid, or erythroid lineages in a hematopoietic setting.

Stem cells may be characterized by both the presence of markers associated with specific epitopes identified by antibodies and the absence of certain markers as identified by

the lack of binding of specific antibodies. Stem cells may also be identified by functional assays both *in vitro* and *in vivo*, particularly assays relating to the ability of stem cells to give rise to multiple differentiated progeny.

Stem cells of interest include hematopoietic stem cells and progenitor cells derived therefrom (U.S. Pat. No. 5,061,620); neural crest stem cells (see Morrison *et al.* (1999) Cell 96:737-749); embryonic stem cells; mesenchymal stem cells; mesodermal stem cells; *etc.*

Other hematopoietic "progenitor" cells of interest include cells dedicated to lymphoid lineages, e.g. immature T cell and B cell populations. The methods of the present invention are useful in expanding selected populations of these cells.

Purified populations of stem or progenitor cells may be used to initiate the cultures. For example, human hematopoietic stem cells may be positively selected using antibodies specific for CD34, thy-1; or negatively selected using lineage specific markers which may include glycophorin A, CD3, CD24, CD16, CD14, CD38, CD45RA, CD36, CD2, CD19, CD56, CD66a, and CD66b; T cell specific markers, tumor specific markers, *etc.* Markers useful for the separation of mesodermal stem cells include FcγRII, FcγRIII, Thy-1, CD44, VLA-4α, LFA-1β, HSA, ICAM-1, CD45, Aa4.1, Sca-1, *etc.* Neural crest stem cells may be positively selected with antibodies specific for low-affinity nerve growth factor receptor (LNGFR), and negatively selected for the markers sulfatide, glial fibrillary acidic protein (GFAP), myelin protein P₀, peripherin and neurofilament. Human mesenchymal stem cells may be positively separated using the markers SH2, SH3 and SH4.

The cells of interest are typically mammalian, where the term refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, laboratory, sports, or pet animals, such as dogs, horses, cats, cows, mice, rats, rabbits, *etc.* Preferably, the mammal is human.

The cells which are employed may be fresh, frozen, or have been subject to prior culture. They may be fetal, neonate, adult. Hematopoietic cells may be obtained from fetal liver, bone marrow, blood, particularly G-CSF or GM-CSF mobilized peripheral blood, or any other conventional source. The manner in which the stem cells are separated from other cells of the hematopoietic or other lineage is not critical to this invention. As described above, a substantially homogeneous population of stem or progenitor cells may be obtained by selective isolation of cells free of markers associated with differentiated cells, while displaying epitopic characteristics associated with the stem cells.

Culture medium: The stem or progenitor cells are grown *in vitro* in an appropriate liquid nutrient medium. Generally, the seeding level will be at least about 10 cells/ml, more usually at least about 100 cells/ml and generally not more than about 10⁵ cells/ml, usually not more than about 10⁴ cells/ml.

Various media are commercially available and may be used, including Ex vivo serum free medium; Dulbecco's Modified Eagle Medium (DMEM), RPMI, Iscove's medium, etc. The medium may be supplemented with serum or with defined additives. Appropriate antibiotics to prevent bacterial growth and other additives, such as pyruvate (0.1-5 mM), glutamine (0.5-5 mM), 2-mercaptoethanol ($1-10 \times 10^{-5}$ M) may also be included.

Culture in serum-free medium is of particular interest. The medium may be any conventional culture medium, generally supplemented with additives such as iron-saturated transferrin, human serum albumin, soy bean lipids, linoleic acid, cholesterol, alpha thioglycerol, crystalline bovine hemin, etc., that allow for the growth of hematopoietic cells.

Preferably the expansion medium is free of cytokines, particularly cytokines that induce cellular differentiation. The term cytokine may include lymphokines, monokines and growth factors. Included among the cytokines are thrombopoietin (TPO); nerve growth factors such as NGF- β ; platelet-growth factor; transforming growth factors (TGFs) such as TGF- α and TGF- β ; erythropoietin (EPO); interferons such as interferon- α , - β , and - γ ; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1 α , IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-11, IL-12; etc. In some circumstances, proliferative factors that do not induce cellular differentiation may be included in the cultures, e.g. c-kit ligand, LIF, and the like.

EXPANSION OF STEM/PROGENITOR CELLS

A population of cells comprising progenitor and/or stem cells is cultured *in vitro* in the presence of enhanced levels of β -catenin, either by genetically altering the cells, or by providing exogenous β -catenin, as described above. The upregulation in β -catenin is sufficient to maintain or increase the number of assayable progenitor cells in the culture. The number of assayable progenitor cells may be demonstrated by a number of assays. After one week the progenitor cell cloning efficiency will usually be at least about 75% that of the starting cell population, more usually 100% that of the starting cell population, and may be as high as 200% that of the starting cell population.

Following the initial period, there is an increased expansion, where the number of assayable cells having the functional phenotype of the initial cell population can increase from about 5 to about 100 fold or more. After this time, the cells can remain in cycle, and expansion is limited primarily by considerations of space. The cells can be frozen using conventional methods at any time, usually after the first week of culture.

Frequently stem cells are isolated from biological sources in a quiescent state. Certain expression vectors, particularly retroviral vectors, do not effectively infect non-cycling cells. Cultures established with these vectors as a source of beta-catenin sequences are

induced to enter the cell cycle by a short period of time in culture with growth factors. For example, hematopoietic stem cells are induced to divide by culture with c-kit ligand, which may be combined with LIF, IL-11 and thrombopoietin. After 24 to 72 hours in culture with cytokines, the medium is changed, and the cells are contacted with the retroviral culture, using culture conditions as described above.

After seeding the culture medium, the culture medium is maintained under conventional conditions for growth of mammalian cells, generally about 37° C and 5% CO₂ in 100% humidified atmosphere. Fresh media may be conveniently replaced, in part, by removing a portion of the media and replacing it with fresh media. Various commercially available systems have been developed for the growth of mammalian cells to provide for removal of adverse metabolic products, replenishment of nutrients, and maintenance of oxygen. By employing these systems, the medium may be maintained as a continuous medium, so that the concentrations of the various ingredients are maintained relatively constant or within a predescribed range. Such systems can provide for enhanced maintenance and growth of the subject cells using the designated media and additives.

These cells may find various applications for a wide variety of purposes. The cell populations may be used for screening various additives for their effect on growth and the mature differentiation of the cells. In this manner, compounds which are complementary, agonistic, antagonistic or inactive may be screened, determining the effect of the compound in relationship with one or more of the different cytokines.

The populations may be employed as grafts for transplantation. For example, hematopoietic cells are used to treat malignancies, bone marrow failure states and congenital metabolic, immunologic and hematologic disorders. Marrow samples may be taken from patients with cancer, and enriched populations of hematopoietic stem cells isolated by means of density centrifugation, counterflow centrifugal elutriation, monoclonal antibody labeling and fluorescence activated cell sorting. The stem cells in this cell population are then expanded *in vitro* and can serve as a graft for autologous marrow transplantation. The graft will be infused after the patient has received curative chemotherapy.

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is

weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference.

The present invention has been described in terms of particular embodiments found or proposed by the present inventor to comprise preferred modes for the practice of the invention. It will be appreciated by those of skill in the art that, in light of the present disclosure, numerous modifications and changes can be made in the particular embodiments exemplified without departing from the intended scope of the invention. For example, due to codon redundancy, changes can be made in the underlying DNA sequence without affecting the protein sequence. Moreover, due to biological functional equivalency considerations, changes can be made in protein structure without affecting the biological action in kind or amount. All such modifications are intended to be included within the scope of the appended claims.

EXPERIMENTAL

Bone marrow cells from BCL2 transgenic mice were isolated, enriched for c-kit over a magnetic column, and then stained with antibodies to sort the Sca1⁺ Thy1.1^{lo} c-kit⁺ lin^{-lo} population on a cell sorter. The cells were double sorted to ensure a high level of purity.

The cells were cultured to initiate cell cycle with Steel factor 100ng/ml with 5% serum in X-Vivo 15 containing the retrovirus. At 3 days 50% of media was replaced with only X-vivo 15, and this dilution was repeated every 2 days. The cells were then cultured with supernatant containing retrovirus encoding activated beta-catenin and recombinant steel factor. The increased growth of the stem cells is shown in Figure 1.

The retroviral supernatant had been generated in commercially available X-vivo 15 media using phoenix cells and a MSCV retroviral construct containing beta-catenin driven by the LTR. The retroviral construct is called MSCV and contains an IRES-GFP, in order to label infected cells. The activating beta catenin mutation is a mutation at the amino terminus that prevents phosphorylation and subsequent degradation by proteosomes. The accumulation of beta catenin in the cytosol allows it to translocate to the nucleus where it associates with the LEF/TCF family of transcription factors to turn on gene expression.

50% of the culture supernatant was replaced every day for 3 days. At the end of this culture period the media was replaced with X-vivo 15. Clusters of cells grew out of this culture, and were analyzed at 5 weeks. By May-Gruenwald-Geimsa staining, these cells appeared to have an immature phenotype with large nuclei and small cytoplasm. By FACS staining a majority of cells are Thy1^{lo} Sca-1⁺ Lin^{lo}-kit^{lo}, a phenotype resembling that of stem

cells. About 50% of the cells are Lin⁻ (LT-HSC phenotype), and 50% Lin^{lo} (ST-HSC phenotype). The analysis is shown in Figure 2.

These cells give rise to lineage positive cells at 4 weeks when transplanted into lethally irradiated mice suggesting that they are able to differentiate to various lineages *in vivo*, while remaining immature *in vitro*.

Lethally irradiated mice were injected with 300, 000 host bone marrow and 100,000 cultured cells. Peripheral blood was take at a later time, at 2 weeks, 3 weeks and 4 weeks so far. Donor type was marked with Ly5.1+ cells. Level of differentiation was determined by using antibodies to mature lineage markers. The results are shown in Figure 3, demonstrating that stem cells over-expressing β -catenin have the ability to give rise to multiple lineages when transplanted.

WHAT IS CLAIMED IS:

1. A method for *in vitro* expansion of mammalian stem or progenitor cells, the method comprising:

5 increasing the intracellular concentration of β -catenin in a progenitor or stem cell in an *in vitro* culture medium for a period of time sufficient for said progenitor or stem cell to divide;

wherein the number of cells having the functional phenotype of said stem or progenitor cells is expanded.

10 2. The method of Claim 1, wherein said stem or progenitor cell is a stem cell.

3. The method of Claim 2, wherein said stem cell is a hematopoietic stem cell.

4. The method of Claim 2, wherein said stem cell is a neural crest stem cell.

15

5. The method of Claim 2, wherein said stem cell is a mesenchymal stem cell.

6. The method of Claim 2, wherein said stem cell is an embryonic stem cell.

20

7. The method of Claim 3, wherein said hematopoietic stem cell is a human cell.

8. The method of Claim 1, wherein said step of increasing the intracellular concentration of β -catenin comprises:

25 introduction of an exogenous nucleic acid comprising beta-catenin coding sequences operably linked to a promoter.

9. The method of Claim 8, wherein said beta-catenin is a wild-type beta-catenin.

10. The method of Claim 8, wherein said beta-catenin is a stabilized mutant beta-catenin.

30

11. The method of Claim 8, wherein said exogenous nucleic acid is a retroviral vector.

35

12. The method of Claim 11, wherein said retroviral vector comprises sites for recombination, flanking said beta-catenin coding sequences.

13. The method of Claim 8, wherein said exogenous nucleic acid is an episomal vector.

14. The method of Claim 1, wherein said step of increasing the intracellular
5 concentration of β -catenin comprises:
addition of exogenous beta-catenin to said culture medium.

15. The method of Claim 14, wherein said beta-catenin is a wild-type beta-catenin.

10

16. The method of Claim 14, wherein said beta-catenin is a stabilized mutant beta-catenin.

15

17. The method of Claim 14, wherein said beta-catenin is genetically fused to a transport moiety.

18. The method of Claim 17, wherein said transport moiety is a fragment of HIV tat protein.

20

19. The method of Claim 1, wherein said stem or progenitor cell is a progenitor cell.

20. The method of Claim 19, wherein said progenitor cell is a hematopoietic progenitor cell.

25

21. The method of Claim 20, wherein said hematopoietic progenitor cell is a lymphoid cell.

22. The method of Claim 21, wherein said lymphoid cell is a B cell.

30

23. The method of Claim 21, wherein said lymphoid cell is a T cell.




Figure 1

2/3




Figure 2

3/3




Figure 3

SEQUENCE LISTING

<110> Tannishtha Reya
Roeland Nusse
Irving L. Weissman

<120> Use of beta-catenin in the expansion of
stem and progenitor cells

<130> SUN-175WO

<160> 4

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 3362

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (215)...(2560)

<400> 1

```

aagcctctcg gtctgtggca gcagcgttgg cccggccccc ggagcggaga gcgagggggag      60
gcggagacgg aggaaggtct gaggagcagc ttcagtcccc gccgagccgc caccgcaggt      120
cgaggacggg cggactcccc cggcggggagg agcctgttcc cctgagggta tttgaagtat      180
accatacaac tgttttgaaa atccagcgtg gaca atg gct act caa gct gat ttg      235
                               Met Ala Thr Gln Ala Asp Leu
                               1                               5

atg gag ttg gac atg gcc atg gaa cca gac aga aaa gcg gct gtt agt      283
Met Glu Leu Asp Met Ala Met Glu Pro Asp Arg Lys Ala Ala Val Ser
      10                               15                               20

cac tgg cag caa cag tct tac ctg gac tct gga atc cat tct ggt gcc      331
His Trp Gln Gln Gln Ser Tyr Leu Asp Ser Gly Ile His Ser Gly Ala
      25                               30                               35

act acc aca gct cct tct ctg agt ggt aaa ggc aat cct gag gaa gag      379
Thr Thr Thr Ala Pro Ser Leu Ser Gly Lys Gly Asn Pro Glu Glu Glu
      40                               45                               50                               55

gat gtg gat acc tcc caa gtc ctg tat gag tgg gaa cag gga ttt tct      427
Asp Val Asp Thr Ser Gln Val Leu Tyr Glu Trp Glu Gln Gly Phe Ser
                               60                               65                               70

cag tcc ttc act caa gaa caa gta gct gat att gat gga cag tat gca      475
Gln Ser Phe Thr Gln Glu Gln Val Ala Asp Ile Asp Gly Gln Tyr Ala
                               75                               80                               85

atg act cga gct cag agg gta cga gct gct atg ttc cct gag aca tta      523
Met Thr Arg Ala Gln Arg Val Arg Ala Ala Met Phe Pro Glu Thr Leu
      90                               95                               100

gat gag ggc atg cag atc cca tct aca cag ttt gat gct gct cat ccc      571
Asp Glu Gly Met Gln Ile Pro Ser Thr Gln Phe Asp Ala Ala His Pro
      105                               110                               115

```

act aat gtc cag cgt ttg gct gaa cca tca cag atg ctg aaa cat gca	619
Thr Asn Val Gln Arg Leu Ala Glu Pro Ser Gln Met Leu Lys His Ala	
120 125 130 135	
ggt gta aac ttg att aac tat caa gat gat gca gaa ctt gcc aca cgt	667
Val Val Asn Leu Ile Asn Tyr Gln Asp Asp Ala Glu Leu Ala Thr Arg	
140 145 150	
gca atc cct gaa ctg aca aaa ctg cta aat gac gag gac cag gtg gtg	715
Ala Ile Pro Glu Leu Thr Lys Leu Leu Asn Asp Glu Asp Gln Val Val	
155 160 165	
ggt aat aag gct gca gtt atg gtc cat cag ctt tct aaa aag gaa gct	763
Val Asn Lys Ala Ala Val Met Val His Gln Leu Ser Lys Lys Glu Ala	
170 175 180	
tcc aga cac gct atc atg cgt tct cct cag atg gtg tct gct att gta	811
Ser Arg His Ala Ile Met Arg Ser Pro Gln Met Val Ser Ala Ile Val	
185 190 195	
cgt acc atg cag aat aca aat gat gta gaa aca gct cgt tgt acc gct	859
Arg Thr Met Gln Asn Thr Asn Asp Val Glu Thr Ala Arg Cys Thr Ala	
200 205 210 215	
ggg acc ttg cat aac ctt tcc cat cat cgt gag ggc tta ctg gcc atc	907
Gly Thr Leu His Asn Leu Ser His His Arg Glu Gly Leu Leu Ala Ile	
220 225 230	
ttt aag tct gga ggc att cct gcc ctg gtg aaa atg ctt ggt tca cca	955
Phe Lys Ser Gly Gly Ile Pro Ala Leu Val Lys Met Leu Gly Ser Pro	
235 240 245	
gtg gat tct gtg ttg ttt tat gcc att aca act ctc cac aac ctt tta	1003
Val Asp Ser Val Leu Phe Tyr Ala Ile Thr Thr Leu His Asn Leu Leu	
250 255 260	
tta cat caa gaa gga gct aaa atg gca gtg cgt tta gct ggt ggg ctg	1051
Leu His Gln Glu Gly Ala Lys Met Ala Val Arg Leu Ala Gly Gly Leu	
265 270 275	
cag aaa atg gtt gcc ttg ctc aac aaa aca aat gtt aaa ttc ttg gct	1099
Gln Lys Met Val Ala Leu Leu Asn Lys Thr Asn Val Lys Phe Leu Ala	
280 285 290 295	
att acg aca gac tgc ctt caa att tta gct tat ggc aac caa gaa agc	1147
Ile Thr Thr Asp Cys Leu Gln Ile Leu Ala Tyr Gly Asn Gln Glu Ser	
300 305 310	
aag ctc atc ata ctg gct agt ggt gga ccc caa gct tta gta aat ata	1195
Lys Leu Ile Ile Leu Ala Ser Gly Gly Pro Gln Ala Leu Val Asn Ile	
315 320 325	
atg agg acc tat act tac gaa aaa cta ctg tgg acc aca agc aga gtg	1243
Met Arg Thr Tyr Thr Tyr Glu Lys Leu Leu Trp Thr Thr Ser Arg Val	
330 335 340	
ctg aag gtg cta tct gtc tgc tct agt aat aag ccg gct att gta gaa	1291
Leu Lys Val Leu Ser Val Cys Ser Ser Asn Lys Pro Ala Ile Val Glu	
345 350 355	

gct ggt gga atg caa gct tta gga ctt cac ctg aca gat cca agt caa Ala Gly Gly Met Gln Ala Leu Gly Leu His Leu Thr Asp Pro Ser Gln 360 365 370 375	1339
cgt ctt gtt cag aac tgt ctt tgg act ctc agg aat ctt tca gat gct Arg Leu Val Gln Asn Cys Leu Trp Thr Leu Arg Asn Leu Ser Asp Ala 380 385 390	1387
gca act aaa cag gaa ggg atg gaa ggt ctc ctt ggg act ctt gtt cag Ala Thr Lys Gln Glu Gly Met Glu Gly Leu Leu Gly Thr Leu Val Gln 395 400 405	1435
ctt ctg ggt tca gat gat ata aat gtg gtc acc tgt gca gct gga att Leu Leu Gly Ser Asp Asp Ile Asn Val Val Thr Cys Ala Ala Gly Ile 410 415 420	1483
ctt tct aac ctc act tgc aat aat tat aag aac aag atg atg gtc tgc Leu Ser Asn Leu Thr Cys Asn Asn Tyr Lys Asn Lys Met Met Val Cys 425 430 435	1531
caa gtg ggt ggt ata gag gct ctt gtg cgt act gtc ctt cgg gct ggt Gln Val Gly Gly Ile Glu Ala Leu Val Arg Thr Val Leu Arg Ala Gly 440 445 450 455	1579
gac agg gaa gac atc act gag cct gcc atc tgt gct ctt cgt cat ctg Asp Arg Glu Asp Ile Thr Glu Pro Ala Ile Cys Ala Leu Arg His Leu 460 465 470	1627
acc agc cga cac caa gaa gca gag atg gcc cag aat gca gtt cgc ctt Thr Ser Arg His Gln Glu Ala Glu Met Ala Gln Asn Ala Val Arg Leu 475 480 485	1675
cac tat gga cta cca gtt gtg gtt aag ctc tta cac cca cca tcc cac His Tyr Gly Leu Pro Val Val Val Lys Leu Leu His Pro Pro Ser His 490 495 500	1723
tgg cct ctg ata aag gct act gtt gga ttg att cga aat ctt gcc ctt Trp Pro Leu Ile Lys Ala Thr Val Gly Leu Ile Arg Asn Leu Ala Leu 505 510 515	1771
tgt ccc gca aat cat gca cct ttg cgt gag cag ggt gcc att cca cga Cys Pro Ala Asn His Ala Pro Leu Arg Glu Gln Gly Ala Ile Pro Arg 520 525 530 535	1819
cta gtt cag ttg ctt gtt cgt gca cat cag gat acc cag cgc cgt acg Leu Val Gln Leu Leu Val Arg Ala His Gln Asp Thr Gln Arg Arg Thr 540 545 550	1867
tcc atg ggt ggg aca cag cag caa ttt gtg gag ggg gtc cgc atg gaa Ser Met Gly Gly Thr Gln Gln Gln Phe Val Glu Gly Val Arg Met Glu 555 560 565	1915
gaa ata gtt gaa ggt tgt acc gga gcc ctt cac atc cta gct cgg gat Glu Ile Val Glu Gly Cys Thr Gly Ala Leu His Ile Leu Ala Arg Asp 570 575 580	1963
gtt cac aac cga att gtt atc aga gga cta aat acc att cca ttg ttt Val His Asn Arg Ile Val Ile Arg Gly Leu Asn Thr Ile Pro Leu Phe 585 590 595	2011

gtg cag ctg ctt tat tct ccc att gaa aac atc caa aga gta gct gca	2059
Val Gln Leu Leu Tyr Ser Pro Ile Glu Asn Ile Gln Arg Val Ala Ala	
600 605 610 615	
ggg gtc ctc tgt gaa ctt gct cag gac aag gaa gct gca gaa gct att	2107
Gly Val Leu Cys Glu Leu Ala Gln Asp Lys Glu Ala Ala Glu Ala Ile	
620 625 630	
gaa gct gag gga gcc aca gct cct ctg aca gag tta ctt cac tct agg	2155
Glu Ala Glu Gly Ala Thr Ala Pro Leu Thr Glu Leu Leu His Ser Arg	
635 640 645	
aat gaa ggt gtg gcg aca tat gca gct gct gtt ttg ttc cga atg tct	2203
Asn Glu Gly Val Ala Thr Tyr Ala Ala Ala Val Leu Phe Arg Met Ser	
650 655 660	
gag gac aag cca caa gat tac aag aaa cgg ctt tca gtt gag ctg acc	2251
Glu Asp Lys Pro Gln Asp Tyr Lys Lys Arg Leu Ser Val Glu Leu Thr	
665 670 675	
agc tct ctc ttc aga aca gag cca atg gct tgg aat gag act gct gat	2299
Ser Ser Leu Phe Arg Thr Glu Pro Met Ala Trp Asn Glu Thr Ala Asp	
680 685 690 695	
ctt gga ctt gat att ggt gcc cag gga gaa ccc ctt gga tat cgc cag	2347
Leu Gly Leu Asp Ile Gly Ala Gln Gly Glu Pro Leu Gly Tyr Arg Gln	
700 705 710	
gat gat cct agc tat cgt tct ttt cac tct ggt gga tat ggc cag gat	2395
Asp Asp Pro Ser Tyr Arg Ser Phe His Ser Gly Gly Tyr Gly Gln Asp	
715 720 725	
gcc ttg ggt atg gac ccc atg atg gaa cat gag atg ggt ggc cac cac	2443
Ala Leu Gly Met Asp Pro Met Met Glu His Glu Met Gly Gly His His	
730 735 740	
cct ggt gct gac tat cca gtt gat ggg ctg cca gat ctg ggg cat gcc	2491
Pro Gly Ala Asp Tyr Pro Val Asp Gly Leu Pro Asp Leu Gly His Ala	
745 750 755	
cag gac ctc atg gat ggg ctg cct cca ggt gac agc aat cag ctg gcc	2539
Gln Asp Leu Met Asp Gly Leu Pro Pro Gly Asp Ser Asn Gln Leu Ala	
760 765 770 775	
tgg ttt gat act gac ctg taa atcatccttt agctgtattg tctgaacttg	2590
Trp Phe Asp Thr Asp Leu *	
780	
cattgtgatt ggcctgtaga gttgctgaga gggctcgagg ggtgggctgg tatctcagaa	2650
agtgcctgac aactaacca agctgagttt cctatgggaa caattgaagt aaactttttg	2710
ttctggctct ttttggtcga ggagtaacaa tacaaatgga ttttgaggagt gactcaagaa	2770
gtgaagaatg cacaagaatg gatcacaaga tgggaatttag caaaccttag ccttgcttgt	2830
taaaattttt tttttttttt ttttaagaat atctgtaatg gtactgactt tgcttgcttt	2890
gaagtagctc tttttttttt tttttttttt tttttttgca gtaactgttt tttaagtctc	2950
tcgtagtgtt aagttatagt gaatactgct acagcaattt ctaattttta agaattgagt	3010
aatggtgtag aactaataat aattcataat cactctaatt aattgtaatc tgaataaagt	3070
gtaacaattg tgtagccttt ttgtataaaa tagacaaata gaaaatggtc caattagttt	3130
cctttttaat atgcttaaaa taagcaggtg gatctatttc atgtttttga tcaaaaacta	3190
tttgggatat gtatgggtag ggtaaatacag taagaggtgt tatttggaac cttgttttgg	3250

acagttttacc agttgccttt tatcccaaag ttgttgtaac ctgctgtgat acgatgcttc 3310
aagagaaaat gcggttataa aaaatggttc agaattaaac ttttaattca tt 3362

<210> 2
<211> 781
<212> PRT
<213> Homo sapiens

<400> 2
Met Ala Thr Gln Ala Asp Leu Met Glu Leu Asp Met Ala Met Glu Pro
1 5 10 15
Asp Arg Lys Ala Val Ser His Trp Gln Gln Gln Ser Tyr Leu Asp
20 25 30
Ser Gly Ile His Ser Gly Ala Thr Thr Thr Ala Pro Ser Leu Ser Gly
35 40 45
Lys Gly Asn Pro Glu Glu Glu Asp Val Asp Thr Ser Gln Val Leu Tyr
50 55 60
Glu Trp Glu Gln Gly Phe Ser Gln Ser Phe Thr Gln Glu Gln Val Ala
65 70 75 80
Asp Ile Asp Gly Gln Tyr Ala Met Thr Arg Ala Gln Arg Val Arg Ala
85 90 95
Ala Met Phe Pro Glu Thr Leu Asp Glu Gly Met Gln Ile Pro Ser Thr
100 105 110
Gln Phe Asp Ala Ala His Pro Thr Asn Val Gln Arg Leu Ala Glu Pro
115 120 125
Ser Gln Met Leu Lys His Ala Val Val Asn Leu Ile Asn Tyr Gln Asp
130 135 140
Asp Ala Glu Leu Ala Thr Arg Ala Ile Pro Glu Leu Thr Lys Leu Leu
145 150 155 160
Asn Asp Glu Asp Gln Val Val Val Asn Lys Ala Ala Val Met Val His
165 170 175
Gln Leu Ser Lys Lys Glu Ala Ser Arg His Ala Ile Met Arg Ser Pro
180 185 190
Gln Met Val Ser Ala Ile Val Arg Thr Met Gln Asn Thr Asn Asp Val
195 200 205
Glu Thr Ala Arg Cys Thr Ala Gly Thr Leu His Asn Leu Ser His His
210 215 220
Arg Glu Gly Leu Leu Ala Ile Phe Lys Ser Gly Gly Ile Pro Ala Leu
225 230 235 240
Val Lys Met Leu Gly Ser Pro Val Asp Ser Val Leu Phe Tyr Ala Ile
245 250 255
Thr Thr Leu His Asn Leu Leu Leu His Gln Glu Gly Ala Lys Met Ala
260 265 270
Val Arg Leu Ala Gly Gly Leu Gln Lys Met Val Ala Leu Leu Asn Lys
275 280 285
Thr Asn Val Lys Phe Leu Ala Ile Thr Thr Asp Cys Leu Gln Ile Leu
290 295 300
Ala Tyr Gly Asn Gln Glu Ser Lys Leu Ile Ile Leu Ala Ser Gly Gly
305 310 315 320
Pro Gln Ala Leu Val Asn Ile Met Arg Thr Tyr Thr Tyr Glu Lys Leu
325 330 335
Leu Trp Thr Thr Ser Arg Val Leu Lys Val Leu Ser Val Cys Ser Ser
340 345 350
Asn Lys Pro Ala Ile Val Glu Ala Gly Gly Met Gln Ala Leu Gly Leu
355 360 365
His Leu Thr Asp Pro Ser Gln Arg Leu Val Gln Asn Cys Leu Trp Thr
370 375 380
Leu Arg Asn Leu Ser Asp Ala Ala Thr Lys Gln Glu Gly Met Glu Gly
385 390 395 400

```

Leu Leu Gly Thr Leu Val Gln Leu Leu Gly Ser Asp Asp Ile Asn Val
      405      410      415
Val Thr Cys Ala Ala Gly Ile Leu Ser Asn Leu Thr Cys Asn Asn Tyr
      420      425      430
Lys Asn Lys Met Met Val Cys Gln Val Gly Gly Ile Glu Ala Leu Val
      435      440      445
Arg Thr Val Leu Arg Ala Gly Asp Arg Glu Asp Ile Thr Glu Pro Ala
      450      455      460
Ile Cys Ala Leu Arg His Leu Thr Ser Arg His Gln Glu Ala Glu Met
      465      470      475
Ala Gln Asn Ala Val Arg Leu His Tyr Gly Leu Pro Val Val Val Lys
      485      490      495
Leu Leu His Pro Pro Ser His Trp Pro Leu Ile Lys Ala Thr Val Gly
      500      505      510
Leu Ile Arg Asn Leu Ala Leu Cys Pro Ala Asn His Ala Pro Leu Arg
      515      520      525
Glu Gln Gly Ala Ile Pro Arg Leu Val Gln Leu Leu Val Arg Ala His
      530      535      540
Gln Asp Thr Gln Arg Arg Thr Ser Met Gly Gly Thr Gln Gln Gln Phe
      545      550      555
Val Glu Gly Val Arg Met Glu Glu Ile Val Glu Gly Cys Thr Gly Ala
      565      570      575
Leu His Ile Leu Ala Arg Asp Val His Asn Arg Ile Val Ile Arg Gly
      580      585      590
Leu Asn Thr Ile Pro Leu Phe Val Gln Leu Leu Tyr Ser Pro Ile Glu
      595      600      605
Asn Ile Gln Arg Val Ala Ala Gly Val Leu Cys Glu Leu Ala Gln Asp
      610      615      620
Lys Glu Ala Ala Glu Ala Ile Glu Ala Glu Gly Ala Thr Ala Pro Leu
      625      630      635
Thr Glu Leu Leu His Ser Arg Asn Glu Gly Val Ala Thr Tyr Ala Ala
      645      650      655
Ala Val Leu Phe Arg Met Ser Glu Asp Lys Pro Gln Asp Tyr Lys Lys
      660      665      670
Arg Leu Ser Val Glu Leu Thr Ser Ser Leu Phe Arg Thr Glu Pro Met
      675      680      685
Ala Trp Asn Glu Thr Ala Asp Leu Gly Leu Asp Ile Gly Ala Gln Gly
      690      695      700
Glu Pro Leu Gly Tyr Arg Gln Asp Asp Pro Ser Tyr Arg Ser Phe His
      705      710      715
Ser Gly Gly Tyr Gly Gln Asp Ala Leu Gly Met Asp Pro Met Met Glu
      725      730      735
His Glu Met Gly Gly His His Pro Gly Ala Asp Tyr Pro Val Asp Gly
      740      745      750
Leu Pro Asp Leu Gly His Ala Gln Asp Leu Met Asp Gly Leu Pro Pro
      755      760      765
Gly Asp Ser Asn Gln Leu Ala Trp Phe Asp Thr Asp Leu
      770      775      780

```

<210> 3

<211> 2702

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (98)... (2443)

<400> 3

gaattccgag	cgtcagtgc	ggaggccgat	tccgagcggg	cggccgcgag	gtaggtgaag	60
ctcagcgcag	agctgctgtg	acaccgctgc	gtggaca	atg gct act	caa gct gac	115
			Met	Ala Thr	Gln Ala Asp	
			1		5	
ctg atg gag	ttg gac atg	gcc atg gag	ccg gac aga	aaa gct gct	gtc	163
Leu Met Glu	Leu Asp Met	Ala Met Glu	Pro Asp Arg	Lys Ala Ala	Val	
	10		15		20	
agc cac tgg	cag cag cag	tct tac ttg	gat tct gga	atc cat tct	ggt	211
Ser His Trp	Gln Gln Gln	Ser Tyr Leu	Asp Ser Gly	Ile His Ser	Gly	
	25		30		35	
gcc acc acc	aca gct cct	tcc ctg agt	ggc aag ggc	aac cct gag	gaa	259
Ala Thr Thr	Thr Ala Pro	Ser Leu Ser	Gly Lys Gly	Asn Pro Glu	Glu	
	40		45		50	
gaa gat gtt	gac acc tcc	caa gtc ctt	tat gaa tgg	gag caa ggc	ttt	307
Glu Asp Val	Asp Thr Ser	Gln Val Leu	Tyr Glu Trp	Glu Gln Gly	Phe	
	55		60		65	70
tcc cag tcc	ttc acg caa	gag caa gta	gct gat att	gac ggg cag	tat	355
Ser Gln Ser	Phe Thr Gln	Glu Gln Val	Ala Asp Ile	Asp Gly Gln	Tyr	
	75		80		85	
gca atg act	agg gct cag	agg gtc cga	gct gcc atg	ttc cct gag	acg	403
Ala Met Thr	Arg Ala Gln	Arg Val Arg	Ala Ala Met	Phe Pro Glu	Thr	
	90		95		100	
cta gat gag	ggc atg cag	atc cca tcc	acg cag ttt	gac gct gct	cat	451
Leu Asp Glu	Gly Met Gln	Ile Pro Ser	Thr Gln Phe	Asp Ala Ala	His	
	105		110		115	
ccc act aat	gtc cag cgc	ttg gct gaa	cca tca cag	atg ttg aaa	cat	499
Pro Thr Asn	Val Gln Arg	Leu Ala Glu	Pro Ser Gln	Met Leu Lys	His	
	120		125		130	
gca gtt gtc	aat ttg att	aac tat cag	gat gac gcg	gaa ctt gcc	aca	547
Ala Val Val	Asn Leu Ile	Asn Tyr Gln	Asp Asp Ala	Glu Leu Ala	Thr	
	135		140		145	150
cgt gca att	cct gag ctg	aca aaa ctg	cta aac gat	gag gac cag	gtg	595
Arg Ala Ile	Pro Glu Leu	Thr Lys Leu	Leu Asn Asp	Glu Asp Gln	Val	
	155		160		165	
gta gtt aat	aaa gct gct	gtt atg gtc	cat cag ctt	tcc aaa aag	gaa	643
Val Val Asn	Lys Ala Ala	Val Met Val	His Gln Leu	Ser Lys Lys	Glu	
	170		175		180	
gct tcc aga	cat gcc atc	atg cgc tcc	cct cag atg	gtg tct gcc	att	691
Ala Ser Arg	His Ala Ile	Met Arg Ser	Pro Gln Met	Val Ser Ala	Ile	
	185		190		195	
gta cgc acc	atg cag aat	aca aat gat	gta gag aca	gct cgt tgt	act	739
Val Arg Thr	Met Gln Asn	Thr Asn Asp	Val Glu Thr	Ala Arg Cys	Thr	
	200		205		210	
gct ggg acc	ctt cac aac	ctt tct cac	cac cgc gag	ggc ttg ctg	gcc	787
Ala Gly Thr	Leu His Asn	Leu Ser His	His Arg Glu	Gly Leu Leu	Ala	
	215		220		225	230

atc ttt aag tct ggt ggc atc cca gcg ctg gtg aaa atg ctt ggg tca	835
Ile Phe Lys Ser Gly Gly Ile Pro Ala Leu Val Lys Met Leu Gly Ser	
235 240 245	
cca gtg gat tct gta ctg ttc tac gcc atc acg aca ctg cat aat ctc	883
Pro Val Asp Ser Val Leu Phe Tyr Ala Ile Thr Thr Leu His Asn Leu	
250 255 260	
ctg ctc cat cag gaa gga gct aaa atg gca gtg cgc cta gct ggt gga	931
Leu Leu His Gln Glu Gly Ala Lys Met Ala Val Arg Leu Ala Gly Gly	
265 270 275	
ctg cag aaa atg gtt gct ttg ctc aac aaa aca aac gtg aaa ttc ttg	979
Leu Gln Lys Met Val Ala Leu Leu Asn Lys Thr Asn Val Lys Phe Leu	
280 285 290	
gct att aca aca gac tgc ctt cag atc tta gct tat ggc aat caa gag	1027
Ala Ile Thr Thr Asp Cys Leu Gln Ile Leu Ala Tyr Gly Asn Gln Glu	
295 300 305 310	
agc aag ctc atc att ctg gcc agt ggt gga ccc caa gcc tta gta aac	1075
Ser Lys Leu Ile Ile Leu Ala Ser Gly Gly Pro Gln Ala Leu Val Asn	
315 320 325	
ata atg agg acc tac act tat gag aag ctt ctg tgg acc aca agc aga	1123
Ile Met Arg Thr Tyr Thr Tyr Glu Lys Leu Leu Trp Thr Thr Ser Arg	
330 335 340	
gtg ctg aaa gtg ctg tct gtc tgc tct agc aac aag ccg gcc att gta	1171
Val Leu Lys Val Leu Ser Val Cys Ser Ser Asn Lys Pro Ala Ile Val	
345 350 355	
gaa gct ggt ggg atg cag gca ctg ggg ctt cat ctg aca gac cca agt	1219
Glu Ala Gly Gly Met Gln Ala Leu Gly Leu His Leu Thr Asp Pro Ser	
360 365 370	
cag cga ctt gtt caa aac tgt ctt tgg act ctc aga aac ctt tca gat	1267
Gln Arg Leu Val Gln Asn Cys Leu Trp Thr Leu Arg Asn Leu Ser Asp	
375 380 385 390	
gca gcg act aag cag gaa ggg atg gaa ggc ctc ctt ggg act cta gtg	1315
Ala Ala Thr Lys Gln Glu Gly Met Glu Gly Leu Leu Gly Thr Leu Val	
395 400 405	
cag ctt ctg ggt tcc gat gat ata aat gtg gtc acc tgt gca gct gga	1363
Gln Leu Leu Gly Ser Asp Asp Ile Asn Val Val Thr Cys Ala Ala Gly	
410 415 420	
att ctc tct aac ctc act tgc aat aat tac aaa aac aag atg atg gtg	1411
Ile Leu Ser Asn Leu Thr Cys Asn Asn Tyr Lys Asn Lys Met Met Val	
425 430 435	
tgc caa gtg ggt ggc ata gag gct ctt gta cgc acc gtc ctt cgt gct	1459
Cys Gln Val Gly Gly Ile Glu Ala Leu Val Arg Thr Val Leu Arg Ala	
440 445 450	
ggt gac agg gaa gac atc act gag cct gcc atc tgt gct ctt cgt cat	1507
Gly Asp Arg Glu Asp Ile Thr Glu Pro Ala Ile Cys Ala Leu Arg His	
455 460 465 470	

ctg acc agc cgg cat cag gaa gcc gag atg gcc cag aat gcc gtt cgc Leu Thr Ser Arg His Gln Glu Ala Glu Met Ala Gln Asn Ala Val Arg 475 480 485	1555
ctt cat tat gga ctg cct gtt gtg gtt aaa ctc ctg cac cca cca tcc Leu His Tyr Gly Leu Pro Val Val Val Lys Leu Leu His Pro Pro Ser 490 495 500	1603
cac tgg cct ctg ata aag gca act gtt gga ttg att cga aac ctt gcc His Trp Pro Leu Ile Lys Ala Thr Val Gly Leu Ile Arg Asn Leu Ala 505 510 515	1651
ctt tgc cca gca aat cat gcg cct ttg cgg gaa cag ggt gct att cca Leu Cys Pro Ala Asn His Ala Pro Leu Arg Glu Gln Gly Ala Ile Pro 520 525 530	1699
cga cta gtt cag ctg ctt gta cga gca cat cag gac acc caa cgg cgc Arg Leu Val Gln Leu Leu Val Arg Ala His Gln Asp Thr Gln Arg Arg 535 540 545 550	1747
acc tcc atg ggt gga acg cag cag cag ttt gtg gag ggc gtg cgc atg Thr Ser Met Gly Gly Thr Gln Gln Gln Phe Val Glu Gly Val Arg Met 555 560 565	1795
gag gaa ata gtc gaa ggg tgt act gga gct ctc cac atc ctt gct cgg Glu Glu Ile Val Glu Gly Cys Thr Gly Ala Leu His Ile Leu Ala Arg 570 575 580	1843
gac gtt cac aac cgg att gta atc cga gga ctc aat acc att cca ttg Asp Val His Asn Arg Ile Val Ile Arg Gly Leu Asn Thr Ile Pro Leu 585 590 595	1891
ttt gtg cag ttg ctt tat tct ccc att gaa aat atc caa aga gta gct Phe Val Gln Leu Leu Tyr Ser Pro Ile Glu Asn Ile Gln Arg Val Ala 600 605 610	1939
gca ggg gtc ctc tgt gaa ctt gct cag gac aag gag gct gca gag gcc Ala Gly Val Leu Cys Glu Leu Ala Gln Asp Lys Glu Ala Ala Glu Ala 615 620 625 630	1987
att gaa gct gag gga gcc aca gct ccc ctg aca gag tta ctc cac tcc Ile Glu Ala Glu Gly Ala Thr Ala Pro Leu Thr Glu Leu Leu His Ser 635 640 645	2035
agg aat gaa ggc gtg gca aca tac gca gct gct gtc cta ttc cga atg Arg Asn Glu Gly Val Ala Thr Tyr Ala Ala Ala Val Leu Phe Arg Met 650 655 660	2083
tct gag gac aag cca cag gat tac aag aag cgg ctt tca gtc gag ctg Ser Glu Asp Lys Pro Gln Asp Tyr Lys Lys Arg Leu Ser Val Glu Leu 665 670 675	2131
acc agt tcc ctc ttc agg aca gag cca atg gct tgg aat gag act gca Thr Ser Ser Leu Phe Arg Thr Glu Pro Met Ala Trp Asn Glu Thr Ala 680 685 690	2179
gat ctt gga ctg gac att ggt gcc cag gga gaa gcc ctt gga tat cgc Asp Leu Gly Leu Asp Ile Gly Ala Gln Gly Glu Ala Leu Gly Tyr Arg 695 700 705 710	2227

cag gat gat ccc agc tac cgt tct ttt cac tct ggt gga tac ggc cag 2275
 Gln Asp Asp Pro Ser Tyr Arg Ser Phe His Ser Gly Gly Tyr Gly Gln
 715 720 725
 gat gcc ttg ggg atg gac cct atg atg gag cat gag atg ggt ggc cac 2323
 Asp Ala Leu Gly Met Asp Pro Met Met Glu His Glu Met Gly Gly His
 730 735 740
 cac cct ggt gct gac tat cca gtt gat ggg ctg cct gat ctg gga cac 2371
 His Pro Gly Ala Asp Tyr Pro Val Asp Gly Leu Pro Asp Leu Gly His
 745 750 755
 gcc cag gac ctc atg gat ggg ctg ccc cca ggt gat agc aat cag ctg 2419
 Ala Gln Asp Leu Met Asp Gly Leu Pro Pro Gly Asp Ser Asn Gln Leu
 760 765 770
 gcc tgg ttt gat act gac ctg taa atcgtcctta gtaagaaagc ttataaaaagc 2473
 Ala Trp Phe Asp Thr Asp Leu *
 775 780
 cagtgtgggt gaatacttac tctgcctgca gaactccaga aagacttggt aggggtgggaa 2533
 tggtttttagg cctgttttgta aatctgccac caaacagata cataccttgg aaggagatgt 2593
 tcatgtgtgg aagtttctca cgttgatggt ttgcccacag cttttgcagc gttatactca 2653
 gatgagtaac atttgctggt ttcaacatta atagcagcct tctctctat 2702

<210> 4

<211> 781

<212> PRT

<213> Mus musculus

<400> 4

Met Ala Thr Gln Ala Asp Leu Met Glu Leu Asp Met Ala Met Glu Pro
 1 5 10 15
 Asp Arg Lys Ala Ala Val Ser His Trp Gln Gln Gln Ser Tyr Leu Asp
 20 25 30
 Ser Gly Ile His Ser Gly Ala Thr Thr Ala Pro Ser Leu Ser Gly
 35 40 45
 Lys Gly Asn Pro Glu Glu Glu Asp Val Asp Thr Ser Gln Val Leu Tyr
 50 55 60
 Glu Trp Glu Gln Gly Phe Ser Gln Ser Phe Thr Gln Glu Gln Val Ala
 65 70 75 80
 Asp Ile Asp Gly Gln Tyr Ala Met Thr Arg Ala Gln Arg Val Arg Ala
 85 90 95
 Ala Met Phe Pro Glu Thr Leu Asp Glu Gly Met Gln Ile Pro Ser Thr
 100 105 110
 Gln Phe Asp Ala Ala His Pro Thr Asn Val Gln Arg Leu Ala Glu Pro
 115 120 125
 Ser Gln Met Leu Lys His Ala Val Val Asn Leu Ile Asn Tyr Gln Asp
 130 135 140
 Asp Ala Glu Leu Ala Thr Arg Ala Ile Pro Glu Leu Thr Lys Leu Leu
 145 150 155 160
 Asn Asp Glu Asp Gln Val Val Val Asn Lys Ala Ala Val Met Val His
 165 170 175
 Gln Leu Ser Lys Lys Glu Ala Ser Arg His Ala Ile Met Arg Ser Pro
 180 185 190
 Gln Met Val Ser Ala Ile Val Arg Thr Met Gln Asn Thr Asn Asp Val
 195 200 205
 Glu Thr Ala Arg Cys Thr Ala Gly Thr Leu His Asn Leu Ser His His
 210 215 220

Arg Glu Gly Leu Leu Ala Ile Phe Lys Ser Gly Gly Ile Pro Ala Leu
 225 230 235 240
 Val Lys Met Leu Gly Ser Pro Val Asp Ser Val Leu Phe Tyr Ala Ile
 245 250 255
 Thr Thr Leu His Asn Leu Leu Leu His Gln Glu Gly Ala Lys Met Ala
 260 265 270
 Val Arg Leu Ala Gly Gly Leu Gln Lys Met Val Ala Leu Leu Asn Lys
 275 280 285
 Thr Asn Val Lys Phe Leu Ala Ile Thr Thr Asp Cys Leu Gln Ile Leu
 290 295 300
 Ala Tyr Gly Asn Gln Glu Ser Lys Leu Ile Ile Leu Ala Ser Gly Gly
 305 310 315 320
 Pro Gln Ala Leu Val Asn Ile Met Arg Thr Tyr Thr Tyr Glu Lys Leu
 325 330 335
 Leu Trp Thr Thr Ser Arg Val Leu Lys Val Leu Ser Val Cys Ser Ser
 340 345 350
 Asn Lys Pro Ala Ile Val Glu Ala Gly Gly Met Gln Ala Leu Gly Leu
 355 360 365
 His Leu Thr Asp Pro Ser Gln Arg Leu Val Gln Asn Cys Leu Trp Thr
 370 375 380
 Leu Arg Asn Leu Ser Asp Ala Ala Thr Lys Gln Glu Gly Met Glu Gly
 385 390 395 400
 Leu Leu Gly Thr Leu Val Gln Leu Leu Gly Ser Asp Asp Ile Asn Val
 405 410 415
 Val Thr Cys Ala Ala Gly Ile Leu Ser Asn Leu Thr Cys Asn Asn Tyr
 420 425 430
 Lys Asn Lys Met Met Val Cys Gln Val Gly Gly Ile Glu Ala Leu Val
 435 440 445
 Arg Thr Val Leu Arg Ala Gly Asp Arg Glu Asp Ile Thr Glu Pro Ala
 450 455 460
 Ile Cys Ala Leu Arg His Leu Thr Ser Arg His Gln Glu Ala Glu Met
 465 470 475 480
 Ala Gln Asn Ala Val Arg Leu His Tyr Gly Leu Pro Val Val Val Lys
 485 490 495
 Leu Leu His Pro Pro Ser His Trp Pro Leu Ile Lys Ala Thr Val Gly
 500 505 510
 Leu Ile Arg Asn Leu Ala Leu Cys Pro Ala Asn His Ala Pro Leu Arg
 515 520 525
 Glu Gln Gly Ala Ile Pro Arg Leu Val Gln Leu Leu Val Arg Ala His
 530 535 540
 Gln Asp Thr Gln Arg Arg Thr Ser Met Gly Gly Thr Gln Gln Gln Phe
 545 550 555 560
 Val Glu Gly Val Arg Met Glu Glu Ile Val Glu Gly Cys Thr Gly Ala
 565 570 575
 Leu His Ile Leu Ala Arg Asp Val His Asn Arg Ile Val Ile Arg Gly
 580 585 590
 Leu Asn Thr Ile Pro Leu Phe Val Gln Leu Leu Tyr Ser Pro Ile Glu
 595 600 605
 Asn Ile Gln Arg Val Ala Ala Gly Val Leu Cys Glu Leu Ala Gln Asp
 610 615 620
 Lys Glu Ala Ala Glu Ala Ile Glu Ala Glu Gly Ala Thr Ala Pro Leu
 625 630 635 640
 Thr Glu Leu Leu His Ser Arg Asn Glu Gly Val Ala Thr Tyr Ala Ala
 645 650 655
 Ala Val Leu Phe Arg Met Ser Glu Asp Lys Pro Gln Asp Tyr Lys Lys
 660 665 670
 Arg Leu Ser Val Glu Leu Thr Ser Ser Leu Phe Arg Thr Glu Pro Met
 675 680 685
 Ala Trp Asn Glu Thr Ala Asp Leu Gly Leu Asp Ile Gly Ala Gln Gly
 690 695 700

Glu	Ala	Leu	Gly	Tyr	Arg	Gln	Asp	Asp	Pro	Ser	Tyr	Arg	Ser	Phe	His
705					710					715					720
Ser	Gly	Gly	Tyr	Gly	Gln	Asp	Leu	Gly	Met	Asp	Pro	Met	Met	Glu	
			725					730						735	
His	Glu	Met	Gly	Gly	His	His	Pro	Gly	Ala	Asp	Tyr	Pro	Val	Asp	Gly
			740					745					750		
Leu	Pro	Asp	Leu	Gly	His	Ala	Gln	Asp	Leu	Met	Asp	Gly	Leu	Pro	Pro
		755					760					765			
Gly	Asp	Ser	Asn	Gln	Leu	Ala	Trp	Phe	Asp	Thr	Asp	Leu			
	770					775						780			

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/01459

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 1/02; C12N 5/00, 5/02, 7/00, 15/63, 15/86; C12P 21/04, 21/06
US CL : 435/2, 69.1, 70.1, 455, 235.1, 325, 375, 377

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
U.S. : 435/2, 69.1, 70.1, 455, 235.1, 325, 375, 377

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
MEDLINE, EAST, DIALOG

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DAMALAS et al. Excess beta-catenin promotes accumulation of transcriptionally active p53. EMBO J. 1999, Vol 18. No. 11, pages 3054-3063, see entire document.	1, 8-9, 11-12
Y	CHU et al. Retrovirus-mediated gene transfer into human hematopoietic stem cells. J. Mol. Med. 1998, Vol 76, pages 184-192, see entire document.	1-3, 7-8, 11-12, 19-20
Y	ZARRIN et al. Comparison of CMV, RSV, SV40 viral and V-lambda-1 cellular promoters in B and T lymphoid and non-lymphoid cell lines. Biochimica et Biophysica Acta 1999, Vol. 1446, pages 135-139, see entire document and Figure 2.	1, 19-23
Y	MORIN et al. Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC. Science 1997, Vol. 275, pages 1787-1790, especially page 1789.	9-10
Y	SATOH et al. Successful transfer of ADA gene in vitro into human peripheral blood CD34 positive cells by transfecting EBV-based episomal vectors. FEBS Letters 1998, Vol 441. No.1, pages 39-42, see entire document.	1, 8, 13, 19-23
A	FAGOTTO et al. Cell contact-dependent signaling. Dev. Biol. 1996, Vol 180, pages 445-454, especially pages 449 and 451.	1-3, 7-13, 19-23
A	WILLERT et al. Beta-catenin: a key mediator of Wnt signaling. Curr. Biol. 1998, Vol 8, pages 95-102, see entire document.	1-3, 7-13, 19-23



Further documents are listed in the continuation of Box C.



See patent family annex.

*

Special categories of cited documents:

"A"

document defining the general state of the art which is not considered to be of particular relevance

"E"

earlier application or patent published on or after the international filing date

"L"

documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O"

documents referring to an oral disclosure, use, exhibition or other means

"P"

document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&"

document member of the same patent family

Date of the actual completion of the international search

04 April 2001 (04.04.2001)

Date of mailing of the international search report

27 APR 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703)305-3230

Authorized officer

Bridget E. Bunner

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/Us01/01459

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ZIEGLER et al. Expansion of stem and progenitor cells. Curr. Opin. Hematol. 1998, Vol 5, pages 434-440, see entire document.	1-3, 7-13, 19-23
A	US 5,851,984A (MATTHEWS et al) 22 December 1998 (22.12.1998), see entire document.	1-3, 7-13, 19-23

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PC1/US01/01459

Continuation of Item 4 of the first sheet: The current title is too long under PCT Rule 4.3. The new title suggestion: "Expansion of Stem and Progenitor Cells by beta-catenin".

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-13 and 19-23, in part, drawn to a method for *in vitro* expansion of mammalian stem or progenitor cells comprising increasing intracellular concentration of β -catenin by introduction of an exogenous nucleic acid comprising β -catenin coding sequences.

Group II, claim(s) 1-7 and 14-23, in part, drawn to a method for *in vitro* expansion of mammalian stem or progenitor cells comprising increasing intracellular concentration of β -catenin by addition of exogenous β -catenin.

The inventions listed as Groups I-II do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Groups I-II claim different methods. Group I recites the special technical feature of *in vitro* expansion of stem or progenitor cells comprising introduction of an exogenous beta-catenin nucleic acid molecule which is not required by the method of Groups II. Group II recites the special technical feature of *in vitro* expansion of stem or progenitor cells comprising introduction of an exogenous beta-catenin protein which is not required by the method of Group I.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

- A. hematopoietic stem cell
- B. neural crest stem cell
- C. mesenchymal stem cell
- D. embryonic stem cell

The following claim(s) are generic: 8-23.

The species listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:

The special technical feature of (A) is a hematopoietic stem cell population. This special technical feature is not shared by any of the other species.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack unity of invention because they are not so linked as to form a single general inventive concept under PCT Rule 13.1.

In order for more than one species to be examined, the appropriate additional examination fees must be paid. The species are as follows:

- E. wild-type beta-catenin
- F. stabilized mutant beta-catenin

The following claim(s) are generic: 2-7, 11-13, and 17-23.

The species listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:

The special technical feature of (E) is an isolated beta catenin protein that has not been mutated. This special technical feature is not shared by any of the other species.

Form PCT/ISA/210 (extra sheet) (July 1998)

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)